


TIALAC

FORMULA

Counteracting Lactose Intolerance

counteracts lactose intolerance

it promotes lactose digestion and therefore balances intestinal flora



Dietary Supplement

60 capsules



Lactose Intolerance


It is a typical syndrome of gastrointestinal complaints that occur after the consumption of foods containing lactose.

CAUSE



- The cells of the small intestine fail to produce lactase, resulting to lactase deficiency. Lactase is an enzyme that converts lactose into **glucose** and **galactose**, (digestible forms).

SYMPTOMS

- **FLATULENCE**
- Bloat
- Gasses
- Abdominal pain
- Diarrhea





Mechanisms of Action



Significant decrease ($p < 0,01$) of H₂ levels exhaled (breath test) by administering two different doses of β -galactosidase as opposed to placebo to patients with malabsorption of lactose (without symptoms) or intolerance with symptoms (6).

Reduction of symptoms




Significant decrease of bloat ($n < 0,01$) and abdominal pain ($n < 0,05$) by administering two different doses of β - galactosidase as opposed to placebo.

Lactobacillus Reuteri, Sporogenes, Subtilis

- **Probiotics: microorganisms that offer balance to intestinal flora.**
- **Convert lactose into galactic acid creating lactase.**
- **Regulate intestinal function.**

Therapeutic schemes (β - galactosidase L. Reuteri) are useful for the treatment of lactose intolerance, since they significantly improve the symptoms in comparison to placebo.



COMPOSITION IN ACTIVE INGREDIENTS	Per daily dose (2-4 capsules)
Lactase 100000 IU	90,00 mg
Lactobacilli Reuteri 100 billion CFU/g	1,00 mg 100 $\delta\sigma$. ufc
Spores of Lactobacilli Sporogenes 15 bn CFU/g	33,50 mg 500 $\delta\sigma$. ufc
Spores of Lactobacilli Subtilis 25 bn CFU/g	83,28 mg 2,5 $\delta\sigma$. ufc
Prebiotic Maltodextrin	1,000 mg

RECOMMENDED TO THOSE WHO

- have symptoms of lactose intolerance.

DOSAGE

- Take 2-4 capsules on a daily basis (1 capsule for every 100 gr of milk consumed).

PRECAUTIONS • The product should not be taken in the following case: Hypersensitivity to any of the ingredients. • Do not exceed the recommended daily dose. • Dietary supplements should not be used as a substitute of a balanced diet. • Keep away from young children. • This product is not intended for the prevention, cure or treatment of a human disease. • Consult with your doctor if you are pregnant, breast-feeding, taking pharmaceutical treatment or having health problems.

Notification number of the EMEA: 28341 / 8-4-2013 / Medicines Agency (EMEA) does not serve as a distribution permit from the European Medicines Agency (EMEA)